

## 1.3.A.16 Staff Use of Naloxone (Narcan)

### I Policy Index:



**Date Signed:** 09/07/2018  
**Distribution:** Public  
**Replaces Policy:** N/A  
**Supersedes Policy Dated:** 07/27/2018  
**Affected Units:** All Units  
**Effective Date:** 09/07/2018  
**Scheduled Revision Date:** June 2019  
**Revision Number:** 2  
**Office of Primary Responsibility:** DOC Administration

### II Policy:

The Department of Corrections (DOC) shall establish guidelines and policy governing the issuance, storage and utilization of Naloxone by DOC staff. Naloxone will be made available to designated units of the DOC to treat opioid overdoses and harmful exposure to an opioid by staff or others.

### III Definitions:

#### **Fentanyl:**

A potent synthetic opioid drug. Schedule II Narcotic. Approximately 100 times more potent than morphine and 50 times more potent than heroin as an analgesic. Can be injected, snorted/sniffed, smoked, taken orally by pill or tablet, or spiked onto blotter paper. Can be present in a variety of forms (e.g. powder, tablets, capsules, solutions and rocks).

#### **Heroin:**

A highly addictive drug processed from morphine. This is a rapidly acting opioid. Typically used in a powdered form. Heroin is a Schedule I Narcotic with high potential for abuse.

#### **Hydromorphone:**

Belongs in the opioid class of drugs. Has an analgesic potency of two to eight times greater than that of morphine and a rapid onset of action. Typically taken as a tablet, capsule, oral solution or an injectable. This is a Schedule II drug with an accepted medical use as a pain reliever. High potential for abuse and may lead to psychological or physical dependence.

#### **Morphine:**

A non-synthetic Schedule II Narcotic with high potential for abuse. Derived from opium and used for the treatment of pain. Can be injected or taken as an oral solution as immediate and extended release tablets and capsules.

#### **Naloxone:**

Also known as Narcan, is a medication that can reverse the effects of an overdose of opioid drugs. When administered during an overdose, it can block the effects of opioids on the brain and restore breathing within two to eight minutes. Common methods of administering are intranasal, auto injector or intramuscular. Has no potential for abuse.

**Opium:**

Highly addictive non-synthetic narcotic extracted from the poppy plant. Opium poppy is the key source for many narcotics, including morphine, codeine and heroin. Usually found in a liquid, solid or powder form. Can be smoked, intravenously injected or taken in pill form.

**Opioid:**

Class of drugs containing or derived from opium, including but not limited to, heroin and morphine.

**Opioid Antagonist:**

A drug that nullifies in whole or in part, the administration of an opioid. The opioid antagonist for the purpose of this policy is Naloxone.

**Staff Member:**

For the purposes of this policy, a staff member is any person employed by the DOC, full or part time, including an individual under contract assigned to the DOC, an employee of another State agency assigned to the DOC, authorized volunteers and student interns.

**Synthetic Opioids:**

Substances synthesized in a laboratory that act on the same targets in the brain as natural opioids (e.g. morphine and codeine) to produce analgesic (pain relief) effects. Can be in powdered or tablet form. Acetyl fentanyl is a synthetic opioid that is very resistant to the effects of Naloxone.

**IV Procedure(s):****1. Training and Management of Naloxone:**

- A. Designated DOC staff members will receive training regarding Naloxone. Training will include the procedures described in SDCL §§ [34-20A-101](#) and [34-20A-102](#).
- B. Naloxone training will be provided to designated new-hire staff during pre-service training. Staff determined to be at risk of possible exposure to fentanyl or other dangerous opioids are required to complete the training.
- C. A qualified and licensed physician designated by the DOC will prescribe Naloxone for administration by trained DOC staff. A standing order will be issued by the physician authorizing possession and administration of Naloxone by designated and trained staff. Authorization may include protocols and procedures that must be trained and followed by staff members administering Naloxone (See SDCL §§ [34-20A-101](#) and [34-20A-102](#)).
- D. Any staff member trained in the administration of Naloxone, acting under a standing order issued by the licensed physician, may possess and administer Naloxone to any person exhibiting symptoms of opiate overdose or exposure (See SDCL § [34-20A-98](#)).
- E. The Director of Parole, Director of Juvenile Services and Wardens will designate staff members (Naloxone Coordinator) under their authority to coordinate the process of maintaining Naloxone (does not include Naloxone issued to Health Services), within their unit of responsibility. Responsibilities of the Naloxone Coordinator include:
  1. Assuring the supply and integrity (including expiration dates) of the Naloxone kits issued to the unit and authorized unit staff. Coordinators are responsible for knowing how many Naloxone kits each unit maintains (includes those kits issued to staff and those available in a DOC facility or community office).

2. Assuring the maintenance of records documenting the replacement of any Naloxone damaged, unusable, expired or administered. All Naloxone records and inventory shall be kept current.
3. Assuring designated staff assigned to the unit have received all required training prior to being granted authority to access, possess or administer Naloxone, including any required remedial training.
4. Assuring any administration of Naloxone by staff (does not include Health Services staff) is documented in an Incident Report, case note or other approve report. Coordinators shall maintain current records for their assigned unit documenting any and all staff administration of Naloxone by calendar year.

## 2. Protocol:

- A. Staff responding to a possible opioid overdose or exposure will first make sure the scene is safe.
- B. Staff will maintain universal precautions. Personal protective equipment is effective in protecting from exposure to fentanyl or other opioids that may be present at the scene or on the victim/person.
  1. Always wear gloves when responding to a possible overdose or exposure incident.
  2. Wear eye protection and a properly-fitted mask or respirator, if available.
  3. Avoid actions that may cause any powder to become airborne. Inhalation of airborne powder is most likely to lead to harmful effects, but, is less likely to occur than contact through the skin.
- C. Staff will perform an assessment of the person, which shall include determining responsiveness, breathing and pulse.
- D. If the incident occurs within a DOC facility, the Incident Command System (ICS) will be initiated. Correctional Health staff will be notified immediately.
- E. If the incident occurs within the community, staff encountering the person will ensure local Emergency Services (911) is contacted.
- F. Signs of an opioid emergency caused by overdose or exposure, may include all or some of the following symptoms:
  1. Unusual sleepiness, stupor or coma.
  2. Breathing problems; slow or shallow breathing or respiratory failure.
  3. Constricted or pinpoint pupils.
- G. Staff will determine to the best of their ability whether the person is experiencing an opioid overdose or exposure. Staff will respond and offer assistance consistent with DOC training and to the best of their abilities, provided it is safe to do so. If the person is suspected of suffering from the effects of an opioid overdose or exposure, staff will respond as follows:
  1. If the person is conscious or easily roused, do not give Naloxone.
  2. If the person is NOT conscious with abnormal breathing and a pulse, lay the person on his/her back, tilt the person's head back and provide support to his/her neck. Apply one dose of Naloxone in one nostril. Administer Naloxone as quickly as possible as prolonged respiratory depression may result in damage to the central nervous system or death. Turn the person on his/her side after administering the Naloxone.

3. If the person is NOT conscious with abnormal breathing and NO pulse, CPR and AED should be initiated, as per accepted protocol (rescue breathing). Use standard basic life support safety precautions (e.g. pocket mask, gloves) to address exposure risk. Apply one dose of Naloxone in one nostril. Administer Naloxone as quickly as possible as prolonged respiratory depression may result in damage to the person's central nervous system or lead to death.
    - a. When applying Naloxone, lay the person on his/her back, tilt the head back while providing support to the neck and apply Naloxone to the person's nostril.
    - b. CPR should be continued as deemed necessary.
    - c. If breathing improves, place the person on their side and re-assess frequently.
  4. Continue to monitor the person. Do not stop assistance to the person or leave the person alone, unless directed to do so by responding medical staff or the scene becomes unsafe and staff is at risk of death or serious bodily injury.
  5. If breathing does not improve within two-three minutes, repeat steps in subsections 2 or 3 above using a new kit and administering the Naloxone in the person's other nostril.
  6. Re-administer Naloxone, using a new container, every two to three minutes if the person does not respond or responds and then relapses into respiratory depression. Continue steps in subsections 2 or 3 above if the person's condition remains unchanged until medical staff arrives. Administer Naloxone into alternate nostrils with each dose.
  7. If, at any time, pulses are lost, CPR and AED must be administered as per accepted protocol.
- H. Adolescents and children age five or older should receive the same dose of Naloxone as adults. For infants and children weighing less than 40 pounds, consult with Emergency Medical Services or the 911 operator.
- I. All persons receiving Naloxone must be referred for medical follow-up. Inmates will be seen by Health Services and EMS will be called. The effects of Naloxone may only last for a limited time and the person may experience another opiate emergency as the Naloxone wears off. Persons who have experienced opioid overdose or exposure requiring treatment require evaluation by a medical professional. Administration of Naloxone is not a substitute for medical care.

### 3. Response to Exposure:

- A. If staff have reasonable belief they or others may have been exposed to fentanyl or other dangerous opioid, immediately move away from the source of the possible exposure.
- B. Immediately notify another staff person so they may observe those exposed for signs of exposure (See Section 2 F.)
- C. Advise others responding to the scene of the possible presence of fentanyl or dangerous opioids.
- D. Do not touch eyes, mouth, nose or any skin after touching any potentially contaminated surface or drug.
- E. Wash any exposed skin thoroughly with cool water, and soap if available. **DO NOT use hand sanitizers** as this may enhance absorption through the skin.
- F. If clothing, shoes or personal protective equipment is contaminated or suspected to be contaminated, remove these items and place in a sealed plastic bag or hazardous material bag/container.

#### 4. Storage/Maintenance/Replacement:

- A. Naloxone kits must be carried and stored in a manner consistent with proper storage guidelines recommended by the manufacturer and/or prescriber. Naloxone is sensitive to temperature and sunlight exposure. Steps shall be taken by staff to ensure the integrity of the Naloxone is not compromised by exposure to adverse conditions.
  - 1. Staff who are issued Naloxone shall ensure they have reasonable access to the Naloxone while performing work duties that could cause risk of exposure (e.g. searches, including pat down search of an offender, or being present at an offender's residence, when responding to a possible overdose or exposure incident).
  - 2. To prevent theft, loss, damage, misuse or access by unauthorized persons, Naloxone not carried by staff or otherwise in the immediate physical control of the staff member, should be secured in a safe storage area.
- B. If Naloxone is kept in a staff member's state vehicle, it must be removed from the vehicle and stored indoors when the temperature in the vehicle is expected to be below 59° F or above 86° F. The Naloxone kit must be protected from direct light.
- C. Naloxone kits located in DOC institutions will be kept in a location secure from inmates but accessible to staff trained to administer the Naloxone.
- D. All Naloxone kits will be inspected monthly by the staff member issued the kit or the Naloxone Coordinator.
- E. Used, damaged, missing or expired Naloxone kits shall be reported to the unit's Naloxone Coordinator for documentation. The Coordinator will ensure the kits are replaced in a timely manner.
  - 1. Do not attempt to reuse a Naloxone kit once the Naloxone has been administered. Each container contains a single dose and cannot be reused or partially administered.
- F. Expired Naloxone (per the date specified on the container) or Naloxone damaged by exposure to adverse conditions will be disposed of properly.

#### 5. Immunity from Civil Liability:

- A. The physician issuing the standing order under established rules, pursuant to SDCL § [34-20A-102](#), and the staff member acting under a standing order who administers Naloxone in good faith compliance with the protocols for administering Naloxone, and the DOC, are not civilly liable for injuries that may be the result of administering or attempting to administer Naloxone. The person may not be held to pay damages to any person for injuries or death associated with the administration of Naloxone (See SDCL § [34-20A-103](#)).
- B. A health care professional authorized to prescribe or dispense an opioid antagonist is not subject to any disciplinary action or civil liability for prescribing or dispensing or an opioid antagonist to a person whom the health care professional reasonably believes may be in a position to assist or administer the opioid antagonist to a person at risk for an opioid-related drug overdose (See SDCL § [34-20A-106](#)).
- C. No staff member is liable for any civil damages as a result of their acts of commission or omission arising out of, and in the course of, their rendering in good faith, any emergency care and services during an emergency which is in their judgment indicated and necessary at the time. Such relief from liability for civil damages extends to the operation of any motor vehicle in connection with any such

care or services. Nothing in this section grants any relief to any person causing any damage by his willful, wanton or reckless act of commission or omission (See SDCL § 20-9-4.1).

**V Related Directives:**

SDCL §§ 20-9-4.1, 34-20A-98, 34-20A-100, 34-20A-101, 34-20A-102, 34-20A-103, 34-20A-104, 34-20A-105 and 34-20A-106.

**VI Revision Log:**

**June 2017:** New Policy

**June 2018:** **Added** does not include Naloxone issued to Health Services in Section 1 E. **Added** "Coordinators are responsible for knowing how many Naloxone kits each unit maintains (includes those kits issued to staff and those available in a DOC facility or community office)" in Section 1 E. 1. **Added** "Coordinators shall maintain sufficient records for their assigned unit documenting any and all staff administration of Naloxone by calendar year" in Section 1 E. 4.

**August 2018:** **Revised** policy statement. **Updated** definition of "Fentanyl". **Added** "staff determined to be at risk of possible exposure to fentanyl or other dangerous opioids are required to complete training" in Section 1 B. **Added** "designated by the DOC" and **Added** "trained and" in Section 1 C. **Added** "or exposure" to Section 1 D. **Added** "All naloxone records and inventory shall be kept current" in Section 1 E. 2. **Added** "possess" to Section 1 E. 3. **Added** "case note or other approved report" in Section 1 E. 4. **Added** "or exposure" in Section 2 A. **Added** new B. 1-3 to Section 2. **Added** "constricted" to Section 2 F. 3. **Deleted** 4. in Section 2 F. **Added** "or been exposed to an opioid" and **Added** "Staff will respond and offer assistance consistent with DOC training and to the best of their abilities, provided it is safe to do so" in Section 2 G. **Added** "accepted" and "rescue breathing" and **Added** "use standard basic life support safety precautions (e.g. pocket mask, gloves) to address exposure risk" in Section 2 G. 3. **Added** "scene becomes unsafe and staff is at risk of death or serious bodily injury" in Section 2 G. 4. **Added** "exposure requiring treatment" in Section 2 I. **Added** new Section 3. **Added** 1. And 2. to Section 4. A.

*Denny Kaemingk (original signature on file)*

Denny Kaemingk, Secretary of Corrections

09/07/2018

Date